

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11

Effects of epirubicin, WX-UK1 and epirubicin + WX-UK1
 on BN-472 mammary tumor load in Brown Norway rats




Figure 12

Effects of epirubicin, WX-UK1 and epirubicin + WX-UK1 on BN-472 mammary tumor weight in Brown Norway rats




Figure 13

Effects of epirubicin, WX-UK1 and epirubicin + WX-UK1 administrations on axillary lymph-node weight




Figure 14

Effects of epirubicin, WX-UK1 and epirubicin + WX-UK1
 administrations on intraperitoneal lymph node weight




Figure 15

Effects of epirubicin, WX-UK1 and epirubicin + WX-UK1 administrations on the number of macroscopic lung foci




Figure 16

Effects of paclitaxel, WX-UK1, and paclitaxel + WX-UK1 administrations on axillary lymph-node weight




Figure 17

Effects of paclitaxel, WX-UK1, paclitaxel + WX-UK1
administrations on intraperitoneal lymph-node weight




Figure 18

Effects of paclitaxel, WX-UK1 and paclitaxel + WX-UK1 administrations on the number of macroscopic lung foci




Figure 19

Effects of 5-FU, WX-UK1, and 5-FU + WX-UK1 on BN-472
 mammary tumor growth in Brown Norway rats




Figure 20

Effects of 5-FU, WX-UK1, and 5-FU + WX-UK1 on BN-472 mammary tumor burden in Brown Norway rats




Figure 21

Effects of 5-FU, WX-UK1, and 5-FU + WX-UK1 on BN-472 mammary tumor burden in Brown Norway rats




Figure 22

Anti metastatic effects of 5-FU, WX-UK1, and 5-FU + WX-UK1
 in Brown Norway rats bearing BN-472 mammary tumors




Figure 23

Anti metastatic effects of 5-FU, WX-UK1, and 5-FU + WX-UK1
 in Brown Norway rats bearing BN-472 mammary tumors




Figure 24

Anti metastatic effects of 5-FU, WX-UK1, and 5-FU + WX-UK1
 in Brown Norway rats bearing BN-472 mammary tumors

